


Product Data Sheet

Anti-Human IgG (H+L) Antibodies and conjugates

Cat# 10319	Goat Anti-Human IgG (H+L), unlabeled	Size: 1 mg
Cat# 10320	Goat Anti-Human IgG (H+L)-HRP conjugate	Size: 1 ml
Cat# 10330	Goat Anti-Human IgG (H+L)-FITC conjugate	Size: 0.5 ml
Cat# 10340	Goat Anti-Human IgG (H+L)-Biotin conjugate	Size: 0.5 ml
Cat# 10350	Goat Anti-Human IgG (H+L)-AP conjugate	Size: 1 ml



Immunoglobulin G (IgG)

most common type of antibody found in the circulation.

Immunoglobulin G (IgG) is a type of antibody. It is a protein complex composed of four peptide chains—two identical heavy chains and two identical light chains arranged in a Y-shape typical of antibody monomers. IgG has molecular weight of approximately 150 kDa, heavy or H chain approximately 50 kDa and light or L chain 25 kDa. Each IgG has two antigen binding sites. Representing approximately 75% of serum antibodies in humans, IgG is the

Goats were immunized with antigen grade **human IgG (H+L)**. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography to remove non-specific antibodies. Specificity of antibody has been tested using IEP, immunodiffusion, and ELISA. The product reacts with human IgG with no significant reactivity with other serum proteins. The antibody may recognize other species IgG (H+L or IgA, IgG, IgM etc)). Purified anti-human IgG (H+L) antibody is supplied as unlabeled or conjugated to Biotin, FITC, AP or HRP.

Form and Storage

Cat# 10319, Anti-human IgG (H+L), unlabeled

The antibody is supplied in PBS, pH 7.4, and 0.05% azide in either **lyophilized** (1.0 mg) or **liquid** form (1 mg/ml). Reconstitute powder in PBS in 1.0 ml to prepare 1 mg/ml solution. Store at -20°C in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Cat# 10320, Anti-human IgG (H+L)-HRP-conjugate

Purified antibody was coupled to HRP (RZ>3.0) using periodate method. The molar enzyme to protein (E/P) ratio =4.0. The antibody is supplied in stabilizing buffer, 0.1% proclin-300 as preservative in either **lyophilized** (1 ml) or **liquid** form (0.5-1 mg/ml). Reconstitute powder in PBS in 1 ml. Store at 4°C in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:1,000-1:10,000 ELISA, 1:1K-1:5K for western, and 1:200-1:1000 (IHC).

Cat# 10350, Anti-human IgG (H+L)-AP-conjugate

The conjugate is provided at ~0.5-1 mg/ml as liquid in a stabilizing buffer (50 mM HEPES pH 7.1, 0.1 M NaCl, 1 mM MgCl₂, 0.1 mM ZnCl₂, pH 7.5, containing 0.2% bovine serum albumin, 0.05%

sodium azide). The product should be **stored at 4°C** and is stable for a minimum of 1 year. Do not store diluted solutions.

Suggested conjugate dilutions are 1:1,000-1:10,000 ELISA, 1:1K-1:5K for western, and 1:200-1:1000 (IHC).

Cat# 10330, Anti-human IgG (H+L)-FITC-conjugate

Purified anti-human IgG (H+L) antibody was coupled to FITC at F/P ratio ~3:7. The antibody is supplied in PBS, pH 7.4,

0.2% BSA and 0.05% azide in either **lyophilized** (0.5 ml) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20°C in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:200-1:2000 for immunofluorescence.

Absorption Wavelength: 495 nm

Emission Wavelength: 528 nm

Cat# 10340, Anti-human IgG (H+L)-Biotin-conjugate

Purified anti-human IgG (H+L) antibody was coupled to Biotin using Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) at F/P ratio ~10-20:1. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.5 mg) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20°C in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:5,000-1:30,000 ELISA, 1:2K-1:10K for western.

Recommended Working Dilution for ELISA

Working dilution for the specific application should be determined by the investigator to obtain the best conditions. Working solution should be prepared immediately before use and diluted solution should be discarded.

Related Material available for ADI

Human IgG (H+L), IgG, IgG (H+L), and IgG (H+L) ELISA kits

Anti-human IgG (H+L), IgG, IgG (H+L), IgG (H+L) antibodies and conjugates

ELISA kits for the detection of human Antibodies

Western blot recycling kit (Use the same blot/strip to probe with multiple antibodies) **Strip Antibodies in 5 minutes** (no heating or mercaptoethanol)

10319-Gt-Anti-Hu-conj

160218SV