

Product Data Sheet

Donkey Anti-Rabbit IgG Antibodies and conjugates (preadsorbed)

Cat# 20319-210	Donkey Anti-Rabbit IgG, unlabeled	Size: 200 ug/0.5 ml
	Form: Liquid; Powder	
Cat# 20320-210	Donkey Anti-Rabbit IgG- HRP conjugate	Size: 200 ug/0.5 ml
	Form: Liquid; Powder	
Cat# 20330-210	Donkey Anti-Rabbit IgG - FITC conjugate	Size: 200 ug/0.5 ml
	Form: Liquid; Powder	
Cat# 20340-210	Donkey Anti-Rabbit IgG - Biotin conjugate	Size: 200 ug/0.5 ml
	Form: Liquid; Powder	
Cat# 20350-210	Donkey Anti-Rabbit IgG- Alk. Phos. conjugate	Size: 200 ug/0.5 ml
	Form: Liquid; Powder	

Description

Donkeys were immunized with antigen grade **Rabbit IgG**. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography to remove non-specific antibodies. Specificity of antibody has been tested using IEP, immunodiffusion, and ELISA. Antibodies were preadsorbed with bovine, chicken, goat, G. pig, hamster, horse, mouse, rat, and sheep proteins to minimize reactivity to these species. The product may react with rabbit IgA and IgM with no significant reactivity with other serum proteins. Purified anti-Rabbit IgG (preadsorbed) is supplied as unlabeled or conjugated to Biotin, FITC, AP or HRP.

Form and Storage

Cat# 20319-210, Anti-Rabbit IgG, unlabeled

The antibody is supplied in PBS, pH 7.4, and 0.05% azide in either **lyophilized** (200 ug) or **liquid** form (200 ug/0.5ml). Reconstitute powder in PBS in 0.5 ml to prepare 0.2 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Cat# 20320-210, Anti-Rabbit IgG-HRP-conjugate

Purified antibody was coupled to HRP (RZ>3.0) using periodate method. The molar enzyme to protein (E/P) ratio = 4.0. The antibody is supplied in stabilizing buffer, 0.1% prolcin-300 as preservative in either **lyophilized** (200 ug) or **liquid** form (200 ug/0.5ml). Reconstitute powder in PBS in 0.5 ml. Store at 4oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:1,000-1:10,000 ELISA, 1:1K-1:5K for western, and 1:200-1:1000 (IHC).

Cat# 20330-210, Anti-Rabbit IgG-FITC-conjugate

Purified anti-Rabbit IgG (H+L) antibody was coupled to FITC at F/P ratio ~3:7. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (200 ug) or **liquid** form (200 ug/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 200 ug/0.5 ml stock solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:200-1:2000 for immunofluorescence.

Absorption Wavelength: 495 nm

Emission Wavelength: 528 nm

Cat# 20340-210, Anti-Rabbit IgG-Biotin-conjugate

Purified anti-Rabbit IgG (H+L) antibody was coupled to Biotin using Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) at F/P ratio ~10-20:1. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (200 ug) or **liquid** form (200 ug/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 0.5 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:5,000-1:30,000 ELISA, 1:2K-1:10K for western.

Cat# 20350-210, AP-conjugate

The conjugate is provided at 200 ug/0.5 ml as liquid in a stabilizing buffer (50 mM Tris-150 mM NaCl-1 mM MgCl₂, pH 7.5, containing 1% bovine serum albumin, 0.05% sodium azide and 50% glycerol). The product should be **stored at 4°C** and is stable for a minimum of 1 year. Do not store diluted solutions.

Recommended Working Dilution for ELISA

Working dilution for the specific application should be determined by the investigator to obtain the best conditions. Working solution should be prepared immediately before use and diluted solution should be discarded.

Related Material available for ADI

Rabbit IgG, IgM ELISA kits

Anti-Rabbit IgG (H+L), IgG, IgG (H+L), IgG (H+L) antibodies and conjugates

ELISA kits for the detection of Rabbit Antibodies

Western blot recycling kit (Use the same blot/strip to probe with multiple antibodies) **Strip Antibodies in 5 minutes** (no heating or mercaptoethanol)

20319-20-30-40-50-210

80516A