



Product Data Sheet

Donkey Anti-Goat IgG (H+L) Antibodies and conjugates

Cat# 30234	Donkey Anti-Goat IgG (H+L)- Cy (TRITC) conjugate (adsorbed)	Size: 0.5 ml
Cat# 30235	Donkey Anti-Goat IgG (H+L)- Cy5 conjugate (adsorbed)	Size: 0.5 ml

Description

Donkeys were immunized with antigen grade **Goat IgG (H+L)**. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography to remove non-specific antibodies. Specificity of antibody has been tested using IEP, immunodiffusion, and ELISA. The product reacts with Goat IgG, IgA and IgM with no significant reactivity with other Goat serum proteins. **Antibodies were further adsorbed with serum proteins from several species (Bovine, Chicken, Human, Guinea Pig, Hamster, Horse, Mouse, Rabbit, Rat) to minimize reactivity with these species.** The antibody may recognize other species IgG (H+L or IgA, IgG, IgM etc). Purified anti-Goat IgG (H+L) antibody is supplied as unlabeled or conjugated to Biotin, FITC, Alk. phos., HRP, Rhodamine, phycoerythrin (PE), Texas Red, Cy3, Cy5, and various IRDye (700DX, 800, and 800CW) to accommodate applications in various immunological techniques.

Form and Storage

Cat# 30234, Cy3-conjugate

Purified anti-Goat IgG (H+L) antibody was coupled to **Cy3** (Molecular Weight 949 daltons) at F/P ratio ~9-10:1. The antibody is supplied in PBS, pH 7.4, 0.5% BSA, IgG, and 0.05% azide in either **lyophilized** (0.5 ml) or **liquid** form (0.5 ml at ~0.5 mg/1 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested applications: Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays

Suggested conjugate dilutions are 1:100-1:500 for immunofluorescence. Users must optimize the dilutions for a given technique.

Absorption Wavelength: 550 nm
Emission Wavelength: 570 nm

Cat# 30235, Cy5-conjugate

Purified anti-Goat IgG (H+L) antibody was coupled to **Cy5.29** (Cyanine 5.29-OSu) (Molecular Weight 975 daltons) at F/P ratio ~6:1. The antibody is supplied in PBS, pH 7.4, 0.5% BSA, IgG, and 0.05% azide in either **lyophilized** (0.5 ml) or **liquid** form (0.5 ml at ~0.5 mg/1 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested applications: Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays

Suggested conjugate dilutions are 1:100-1:500 for immunofluorescence. Users must optimize the dilutions for a given technique.

Absorption Wavelength: 650 nm
Emission Wavelength: 667 nm

Recommended Working Dilution for ELISA

Working dilution for the specific application should be determined by the investigator to obtain the best conditions. Working solution should be prepared immediately before use and diluted solution should be discarded.

Origin: All material used in various products are derived from US sources and USDA inspected facilities.

MSDS: A Material Safety Data Sheet is not required for this product. The product does not contain any hazardous components above 1% or any carcinogens above 0.1% as defined in 29 CFR 1910.1200, the OSHA Hazard Communication Standard. This material is not required to appear on the TSCA inventory.

This product is for in vitro research use only.

Related Material available for ADI

Goat IgG, IgM ELISA kits
Anti-Goat IgG (H+L), IgG, IgG (H+L), IgG (H+L) antibodies and conjugates

30234, 30235 71120A