



Product Data Sheet

Anti-Dog IgG (H+L) Antibodies and conjugates

Cat# 30360	Goats Anti-Dog IgG (H+L), unlabeled	Size: 1 mg	Form: Liquid;	Powder
Cat# 30361	Goats Anti-Dog IgG (H+L)-HRP conjugate	Size: 0.5 ml	Form: Liquid;	Powder
Cat# 30362	Goats Anti-Dog IgG (H+L)-Alk. Phos. (AP) conjugate	Size: 0.5 ml	Form: Liquid;	Powder
Cat# 30363	Goats Anti-Dog IgG (H+L)-FITC conjugate	Size: 1 mg	Form: Liquid;	Powder
Cat# 30364	Goats Anti-Dog IgG (H+L)-Biotin conjugate	Size: 0.5 ml	Form: Liquid;	Powder

Description

Goats were immunized with antigen grade **Dog IgG (H+L)**. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography to remove non-specific antibodies. Specificity of antibody has been tested using IEP, immunodiffusion, and ELISA. The product reacts with Dog IgG, IgA and IgM with no significant reactivity with other serum proteins. The antibody may recognize other species IgG (H+L or IgA, IgG, IgM etc). Purified anti-Dog IgG (H+L) antibody is supplied as unlabeled or conjugated to Biotin, FITC, AP or HRP.

BSA and 0.05% azide in either **lyophilized** (1 mg/ml) or **liquid** form (1 mg/ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:200-1:2000 for immunofluorescence.

Absorption Wavelength: 495 nm

Emission Wavelength: 528 nm

Form and Storage

Cat# 30364, Anti-Dog IgG (H+L)-Biotin-conjugate

Cat# 30360, Anti-Dog IgG (H+L), unlabeled

Purified anti-Dog IgG (H+L) antibody was coupled to Biotin using Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) at F/P ratio ~10-20:1. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.5 mg) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

The antibody is supplied in PBS, pH 7.4, and 0.05% azide in either **lyophilized** (1.0 mg) or **liquid** form (1 mg/ml). Reconstitute powder in PBS in 1.0 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:5,000-1:30,000 ELISA, 1:2K-1:10K for western.

Cat# 30361, Anti-Dog IgG (H+L)-HRP-conjugate

Recommended Working Dilution for ELISA

Purified antibody was coupled to HRP (RZ>3.0) using periodate method. The molar enzyme to protein (E/P) ratio = 4.0. The antibody is supplied in stabilizing buffer, 0.1% prolcin-300 as preservative in either **lyophilized** (0.5 ml) or **liquid** form (0.5-1 mg/ml). Reconstitute powder in PBS in 1 ml. Store at 4oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Working dilution for the specific application should be determined by the investigator to obtain the best conditions. Working solution should be prepared immediately before use and diluted solution should be discarded.

Suggested conjugate dilutions are 1:1,000-1:10,000 ELISA, 1:1K-1:5K for western, and 1:200-1:1000 (IHC).

Related Material available for ADI

Cat# 30362, AP-conjugate

Dog IgG,, IgM ELISA kits

The conjugate is provided at ~0.5-1 mg/ml as liquid in a stabilizing buffer (50 mM Tris-150 mM NaCl-1 mM MgCl₂, pH 7.5, containing 1% bovine serum albumin, 0.05% sodium azide and 50% glycerol). The product should be **stored at 4°C** and is stable for a minimum of 1 year. Do not store diluted solutions.

Anti-Dog IgG (H+L), IgG, IgG (H+L), IgG (H+L) antibodies and conjugates

ELISA kits for the detection of Dog Antibodies

Western blot recycling kit (Use the same blot/strip to probe with multiple antibodies) **Strip Antibodies in 5 minutes** (no heating or mercaptoethanol)

Cat# 30363, Anti-Dog IgG (H+L)-FITC-conjugate

30360, 30361, 30362, 30363, 30364

70721A

Purified anti-Dog IgG (H+L) antibody was coupled to FITC at F/P ratio ~3:7. The antibody is supplied in PBS, pH 7.4, 0.2%