



Product Data Sheet

Goat Anti-Chicken IgG (H+L) Antibodies and conjugates

Cat# 60334	Goat Anti-Chicken IgG (H+L)- Cy3 conjugate (adsorbed)	Size: 0.5 ml
Cat# 60335	Goat Anti-Chicken IgG (H+L)- Cy5 conjugate (adsorbed)	Size: 0.5 ml

Description

Goats were immunized with antigen grade **Chicken IgG (H+L)**. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography to remove non-specific antibodies. Specificity of antibody has been tested using IEP, immunodiffusion, and ELISA. The product reacts with Chicken IgG, IgA and IgM with no significant reactivity with other Chicken serum proteins. **Antibodies were further adsorbed with serum proteins from several species (Bovine, Hasmter, G. Pig, Goat, Human, Guinea Pig, Rat, Horse, Rabbit, Mouse, and sheep) to minimize reactivity with these species.** The antibody may recognize other species IgG (H+L or IgA, IgG, IgM etc). Purified anti-Chicken IgG (H+L) antibody is supplied as unlabeled or conjugated to Biotin, FITC, Alk. phos., HRP, Rhodamine, Cy3, and Cy5 to accommodate applications in various immunological techniques.

Form and Storage

Cat# 60334, Cy3-conjugate

Purified anti-Chicken IgG (H+L) antibody was coupled to Cy3 (Molecular Weight 949 daltons) at F/P Chickenio ~9-10:1. The antibody is supplied in PBS, pH 7.4, 0.5% BSA, IgG, and 0.05% azide in either **lyophilized** (0.5 ml) or **liquid** form (0.5 ml at ~0.5 mg/1 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested applications: Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays

Suggested conjugate dilutions are 1:100-1:500 for immunofluorescence. Users must optimize the dilutions for a given technique.

Absorption Wavelength: 550 nm
Emission Wavelength: 570 nm

Cat# 60335, Cy5-conjugate

Purified anti-Chicken IgG (H+L) antibody was coupled to Cy5.29 (Cyanine 5.29-OSu) (Molecular Weight 975 daltons) at F/P Chickenio ~6:1. The antibody is supplied in PBS, pH 7.4, 0.5% BSA, IgG, and 0.05% azide in either **lyophilized** (0.5 ml) or **liquid** form (0.5 ml at ~0.5 mg/1 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested applications: Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays

Suggested conjugate dilutions are 1:100-1:500 for immunofluorescence. Users must optimize the dilutions for a given technique.

Absorption Wavelength: 650 nm
Emission Wavelength: 667 nm

Recommended Working Dilution for ELISA

Working dilution for the specific application should be determined by the investigator to obtain the best conditions. Working solution should be prepared immediately before use and diluted solution should be discarded.

Origin: All material used in various products are derived from US sources and USDA inspected facilities.

MSDS: A Material Safety Data Sheet is not required for this product. The product does not contain any hazardous components above 1% or any carcinogens above 0.1% as defined in 29 CFR 1910.1200, the OSHA Hazard Communication Standard. This material is not required to appear on the TSCA inventory.

This product is for in vitro research use only.

Related Material available for ADI

Chicken IgG, IgM ELISA kits

Anti-Chicken IgG (H+L), IgG, IgG (H+L), IgG (H+L) antibodies and conjugates

60334, 60335 71123A