



Product Data Sheet

Anti-Bovine IgM antibodies and conjugates

Cat# 80719	Rabbit Anti-Bovine IgM, unlabeled	Size: 0.5 mg	Form: <input type="checkbox"/> Liquid <input type="checkbox"/> Powder
Cat# 80720	Rabbit Anti-Bovine IgM-HRP conjugate	Size: 0.5 ml	Form: <input type="checkbox"/> Liquid <input type="checkbox"/> Powder
Cat# 80730	Rabbit Anti-Bovine IgM-FITC conjugate	Size: 0.5 ml	Form: <input type="checkbox"/> Liquid <input type="checkbox"/> Powder
Cat# 80740	Rabbit Anti-Bovine IgM-Biotin conjugate	Size: 0.5 ml	Form: <input type="checkbox"/> Liquid <input type="checkbox"/> Powder

Description

Rabbits were immunized with antigen grade Bovine IgM. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography and solid phase adsorbed to remove reactivity with IgA, IgG and other serum proteins. Purified anti-Bovine IgM is supplied as unlabeled, -HRP, -AP, FITC and biotin conjugate.

Purity/Specificity

Specificity of antibodies has been tested using IEP, immunodiffusion, and ELISA. The antibodies yielded a single precipitin arc against anti-bovine serum and Bovine IgM. No antibody reactivity is detected to other bovine serum proteins, IgA and IgG. Anti-Bovine IgM may crossreact with IgM from other species.

Form and Storage

Cat# 80719, Anti-Bovine IgM, unlabeled

The antibody is supplied in PBS, pH 7.4, and 0.05% azide in either **lyophilized** (0.5 mg) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 0.5 mg/0.5 ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Cat# 80720, Anti-Bovine IgM-HRP-conjugate

Purified antibody was coupled to HRP (RZ>3.0) using periodate method. The molar enzyme to protein (E/P) ratio = 4.0. The antibody is supplied in stabilizing buffer, 0.1% prolcin-300 as preservative in either **lyophilized** (0.5 ml) or **liquid** form (0.5 ml; 0.5-1 mg/ml). Reconstitute powder in PBS in 1 ml. Store at 4oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:1,000-1:10,000 ELISA, 1:1K-1:5K for western, and 1:200-1:1000 (IHC).

Cat# 80730, Anti-Bovine IgM-FITC-conjugate

Purified anti-Bovine IgM antibody was coupled to FITC at F/P ratio ~3:7. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.5 ml) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:200-1:2000 for immunofluorescence.

Absorption Wavelength: 495 nm

Emission Wavelength: 528 nm

Cat# 80740, Anti-Bovine IgM-Biotin-conjugate

Purified anti-Bovine IgM antibody was coupled to Biotin using ADI Biotinylation kit # 80300 at F/P ratio ~10-20:1. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.5 mg) or **liquid** form (0.5 mg/0.5 ml). Reconstitute powder in PBS in 0.5 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:5,000-1:30,000 ELISA, 1:2K-1:10K for western.

Recommended Working Dilution for ELISA

Working dilution for the specific application should be determined by the investigator to obtain the best conditions. Working solution should be prepared immediately before use and diluted solution should be discarded.

Related Material available for ADI

Mouse IgG, IgM, IgA ELISA kits

Anti-Bovine IgM, IgG, IgG (H+L), IgG (H+L) antibodies and conjugates

ELISA kits for the detection of Mouse Antibodies

80719, 80720, 80730, 80740

90206A