

Goat Anti-Horse IgG (H+L) Antibodies and conjugates

Cat# 80819	Goat Anti-Horse IgG (H+L), unlabeled	Size: 1 mg	Form: Liquid;	Powder
Cat# 80820	Goat Anti-Horse IgG (H+L)-HRP conjugate	Size: 1 ml	Form: Liquid;	Powder
Cat# 80830	Goat Anti-Horse IgG (H+L)-FITC conjugate	Size: 1 ml	Form: Liquid;	Powder
Cat# 80840	Goat Anti-Horse IgG (H+L)-Biotin conjugate	Size: 1 ml	Form: Liquid;	Powder
Cat# 80850	Goat Anti-Horse IgG (H+L)-Alk. Phos. conjugate	Size: 0.5 ml	Form: Liquid;	Powder

Description

Goats were immunized with antigen grade **Horse IgG (H+L)**. Antibodies have been isolated using ammonium sulfate, ion-exchange, and affinity chromatography to remove non-specific antibodies. Specificity of antibody has been tested using IEP, immunodiffusion, and ELISA. The product reacts with Horse IgG, and it may also react with IgA and IgM with no significant reactivity with other serum proteins. The antibody may recognize other species IgG (H+L or IgA, IgG, IgM etc)). Purified anti-Horse IgG (H+L) antibody is supplied as unlabeled or conjugated to Biotin, FITC, AP or HRP.

Form and Storage

Cat# 80819, Anti-Horse IgG (H+L), unlabeled

The antibody is supplied in PBS, pH 7.4, and 0.05% azide in either **lyophilized** (1.0 mg) or **liquid** form (1 mg/ml). Reconstitute powder in PBS in 1.0 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Cat# 80820, Anti-Horse IgG (H+L)-HRP-conjugate

Purified antibody was coupled to HRP (RZ>3.0) using periodate method. The molar enzyme to protein (E/P) ratio = 4.0. The antibody is supplied in stabilizing buffer, 0.1% prolcin-300 as preservative in either **lyophilized** (1 ml) or **liquid** form (0.5-1 mg/ml). Reconstitute powder in PBS in 1 ml. Store at 4oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:1,000-1:10,000 ELISA, 1:1K-1:5K for western, and 1:200-1:1000 (IHC).

Cat# 80830, Anti-Horse IgG (H+L)-FITC-conjugate

Purified anti-Horse IgG (H+L) antibody was coupled to FITC at F/P ratio ~3:7. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (1 ml) or **liquid** form (1 ml at 0.5 mg/0.5 ml). Reconstitute powder in PBS in 1 ml to prepare 1 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:200-1:2000 for immunofluorescence.

Absorption Wavelength: 495 nm
Emission Wavelength: 528 nm

Cat# 80840, Anti-Horse IgG (H+L)-Biotin-conjugate

Purified anti-Horse IgG (H+L) antibody was coupled to Biotin using Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) at F/P ratio ~10-20:1. The antibody is supplied in PBS, pH 7.4, 0.2% BSA and 0.05% azide in either **lyophilized** (0.5 mg) or **liquid** form (1 ml at 0.5 mg/0.5 ml). Reconstitute powder in PBS in 1.0 ml to prepare 0.5 mg/ml solution. Store at -20oC in suitable aliquots. Stability is ~6-12 months. Do not freeze and thaw.

Suggested conjugate dilutions are 1:5,000-1:30,000 ELISA, 1:2K-1:10K for western.

Cat# 80850-AP, AP-conjugate

The conjugate is provided at ~0.5-1 mg/ml as liquid in a stabilizing buffer (50 mM Tris-150 mM NaCl-1 mM MgCl₂, pH 7.5, containing 1% bovine serum albumin, 0.05% sodium azide and 50% glycerol). The product should be **stored at 4°C** and is stable for a minimum of 1 year. Do not store diluted solutions.

Recommended Working Dilution for ELISA

Working dilution for the specific application should be determined by the investigator to obtain the best conditions. Working solution should be prepared immediately before use and diluted solution should be discarded.

Related Material available for ADI

Horse IgG, IgM ELISA kits

Anti-Horse IgG (H+L), IgG, IgG (H+L), IgG (H+L) antibodies and conjugates

ELISA kits for the detection of Horse Antibodies

Western blot recycling kit (Use the same blot/strip to probe with multiple antibodies) **Strip Antibodies in 5 minutes** (no heating or mercaptoethanol)

80819, 80820, 80830, 80840, 80850 90922A