



Product Data Sheet

Cat # AB-13110

Mouse Anti-Human CD8-FITC IgG

Size: 100ug

Synonyms:

CD8, MAL, p32

Type:

Mouse Anti Human Monoclonal

Introduction:

CD8 is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. CD8 acts as a co-receptor, and the T-cell receptor on the T lymphocyte recognize antigen displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional CD8 is either a homodimer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. CD8 identifies cytotoxic/suppressor t-cells that interact with MHC class I bearing targets. CD8 is thought to play a role in the process of t-cell mediated killing. CD8 alpha chains binds to class-I MHC molecules alpha-3 domains. Staining antibody, weakly cytotoxic. For staining, use 10 μ l/ 10^6 cells. Titer for cytotoxicity should be determined by the investigator.

Immunogen:

Purified human PBL CD8+ T cells

Ig Subclass:

Mouse IgG2a; purification by ion exchange column; 1mg/mL in PBS after reconstitution; Reconstitute with of H2O. Mix gently, wash the sides of the vial and wait 30-60 seconds before use.

Storage Procedures:

Lyophilized: store at 4°C. After reconstitution, if not intended for use within a month, aliquot and store at -20°C. Two years lyophilized, one month in solution at 4°C.

Available Conjugates:

This antibody is also available conjugated to biotin

Usage:

This item is for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals. If supplied in powder then reconstitute it in 100 μ l water for 1 mg/ml stock and store in liquid at 4°C for ~1 week or aliquots in suitable size and store at -20°C for long term storage..

Rev. 101001V