
Mouse Anti-Human CD7-FITC conjugate

PRODUCT INFORMATION

Catalog # CD07F-100 **SIZE:** 100 Tests

Catalog # CD07F-25 **SIZE:** 25 Tests

CLONE: HIT7

ISOTYPE: Mouse IgG1

WS.No.: □5T CD07.03

Product Forms: Purified, FITC conjugation, PE conjugation.

DESCRIPTION

CD7 McAb recognizes a 40KD type I single-chain transmembrane glycoprotein. CD7 antigen has been considered as one of the earliest and immature T cell markers and is expressed on all thymocytes, most mature T cells, B-lineage progenitors, NK cells, monocytes (weakly) and some stem cell leukemia capable of multilineage differentiation. CD7 antigen may be considered as a marker of undifferentiated hemopoietic cells capable of giving rise to T, B and myeloid lineage cells under the appropriate microenvironment. T cell marker CD7 is also present on the majority cells of T lymphoid leukemia and small party cells of acute myeloid leukemia (AML). Mature B lymphocytes and granulocytes do not express CD7 antigen.

PREPARATION

The monoclonal antibody is purified from ascites by protein G affinity chromatography and is conjugated with FITC, R-PE under optimum conditions.

USAGE

The purified reagent is effective for indirect immunofluorescence staining of human cells for flow cytometric analysis.

The conjugated reagent (FITC, R-PE) is tested for flow cytometric analysis using 20µl/10⁶ cells or 100µl peripheral blood cells.

STORAGE

For purified forms, long term storage at -20oC.

For conjugated forms, storage at 4oC, should not be frozen and avoid prolonged exposure to light.

REFERENCES

Schlossman S., L. Bloumsell, W. Gilks, et al., eds. 1995. Leucocyte Typing □: White Cell Differentiation Antigens. P: 246 Oxford University Press, New York.