

Product Data Sheet

Methoxy polyethylene glycol (mPEG) conjugates

□ Cat # PEG-BSA-20K

mPEG-BSA (Molecular Weight: 20,000)

Size: 100 ug

PEGylation is a process of covalent and non-covalent attachment of polyethylene glycol (PEG) polymer chains to molecules such as drugs or therapeutic proteins. The covalent attachment of PEG to a drug or protein can reduce the immunogenicity, antigenicity, and increase the hydrodynamic size which prolong its circulatory time. PEGylation imparts several pharmacological advantages such as improved drug solubility, reduced dosage frequency, extended circulating life, increased drug stability, and enhanced protection from proteolytic degradation. Currently 15 PEGylated pharmaceuticals are on the market; Adynovate, Plegridy, Peginterferon alfa-2a, Doxorubicin HCL liposome, Peginterferon alfa-2b, Pegaspargase, and Pegademase bovine. ADI's mPEG-BSA conjugates are suitable for use as reference standards or coating antigens in ELISA.



Cat# PEG-BSA-20K: Prepared by conjugation of Methoxy polyethylene glycol to native BSA followed by dialysis to remove free PEG.

Concentration: 1 mg/ml (100 ul) in liquid assuming an absorbance of 0.632 at 280 nm. Refer to vial for lot specific concentration. Please note: The concentration above refers only to the BSA polypeptide concentration. It does not include the mass contributed by the 20 kDa mPEG chains.

Buffer: PBS pH 7.4, 0.05% Sodium Azide

Storage: 4°C for short term (<1 month) and 20°C for long term storage (6-12 months)

Analysis: ELISA & SDS-PAGE



ELISA: 1 ug/ml anti-PEG coated antibody and anti-PEG detection antibody

SDS-PAGE: Lane 1: Ladder. Lane 2: BSA. Lane 3: 2.5 ug PEG-BSA-20K. Due to steric hindrance caused by larger PEG chains the PEG is coupled to BSA at varying degrees causing smears to appear. Some free BSA is detected.

For in vitro Research use only (RUO)

PEG-BSA-20K

010917IA

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, San Antonio, TX 78244, USA; **Email:** service@4adi.com
☎ (800) 786-5777; (210) 561-9515; **Fax:** (210) 561-9544; **Web Site:** www.4adi.com